

Valvular Heart Disease/Cardiac Surgery

Shahbudin H. Rahimtoola, MB, FRCP, MACP, MACC, DSc (Hon)

Los Angeles, California

AORTIC STENOSIS (AS)

Of 5,621 men and women ≥ 65 years of age in the Cardiovascular Health Study (1), 6% progressed from aortic sclerosis to AS over 5 years. Older age, male gender, and shorter height were predictors of progression to AS. Baseline C-reactive protein (CRP) concentration was not a predictor of calcific aortic valve disease at baseline or of its progression.

In a retrospective study of 818 patients (2), native AS progression could not be predicted by abnormalities in the seven markers of atherosclerotic risk.

High-grade AS valves (57 native valves and 24 bioprostheses) showed presence of endothelial progenitor cells, dendritic cells, and inflammatory cell markers (3). These changes were even more frequently found in bioprosthesis; cell density was higher in AS. These data point to a systemic origin of valvular cells and provide "novel insight" into mechanisms of aortic valve stenosis.

One-third of 216 patients age ≥ 75 years were denied aortic valve replacement (AVR) (4). Older age and depressed left ventricular ejection fraction (LVEF) were the characteristics most strongly associated with absence of surgery.

In the I-REVIVE trial (5), 16 patients had percutaneous implantation of an equine pericardial valve. Three patients died early. Among 11 implanted patients, 4 were alive at 6 months with normal valve function. The ReValving system using bovine pericardium for percutaneous valve replacement was initiated (6). A 21-patient feasibility study was also initiated.

MITRAL STENOSIS (MS)

Antibiotic prophylaxis: a meta-analysis was undertaken of 10 trials of young adults living on U.S. military bases from 1950 to 1961 (7). Antibiotic prophylaxis with a single injection of intramuscular (IM) penicillin for group A streptococcal infection was effective for preventing acute rheumatic fever with protective effect of 80% (fixed defect relative risk [RR] 0.20; 95% confidence interval [CI] 0.11 to 0.36); for a need to treat (NTT) number of 60, the marginal cost was \$46.

Catheter balloon commissurotomy (CBC) for MS in 626 patients (8) showed the procedure was effective at a follow-up of 6.5 ± 2.5 years in those age ≥ 41 years, 42 to 53 years, 54 to 63 years, and in those age > 63 years.

In 468 consecutive patients undergoing CBC for MS (9), the 5-, 10-, and 14-year event-free survival was $94 \pm 1\%$,

$76 \pm 9\%$, and $76 \pm 9\%$. Mitral valve score by echocardiogram > 8 and post-procedure mitral valve area (MVA) < 2.0 cm² were predictors of restenosis ($p < 0.0002$ and $p < 0.002$, respectively).

MITRAL REGURGITATION (MR)

Degenerative ($n = 20$), rheumatic ($n = 23$), and control ($n = 20$) mitral valves were studied by electron microscopy and immunochemistry (10). Cellular proliferation and matrix production were shown to be important in the development of mitral valve disease. The severity of calcification correlated with higher levels of cellular proliferation and matrix production in calcified rheumatic valves as compared to degenerative mitral valves.

A total of 132 consecutive asymptomatic patients age 55 ± 15 years who had severe "degenerative" MR were seen from 1995 to 2002 and were followed until 2004. Mean \pm SD for follow-up was 62 ± 26 months (11). There were four deaths: one sudden death in an 80-year-old; three deaths were unrelated to MR. Thirty-two patients had surgery for the following indications: symptoms, LVEF $< 60\%$, left ventricular end-systolic dimension ≥ 45 mm, pulmonary hypertension (pulmonary artery systolic pressure > 50 mm Hg), or new onset of atrial fibrillation (AF). There was no operative mortality. At follow-up, LVEF was normal in 28 of 32 patients, the other 4 patients had mitral valve replacement (MVR) and coronary artery bypass graft surgery (CABG). Survival of the entire group at two years was $99 \pm 1\%$, at four years $96 \pm 2\%$, and at six and eight years was $91 \pm 3\%$.

A retrospective review of the echocardiography database revealed 869 patients who had severe MR and normal LVEF (12). The mean age was 68 ± 17 years, 44% were men, LVEF was $66 \pm 7\%$, diabetes mellitus was present in 19%, hypertension in 65%, and coronary artery disease (CAD) in 39% of patients. Mitral valve surgery was performed in 44%, and 42% of patients were on a beta-blocker. Use of a beta-blocker was associated with a significantly reduced mortality ($p = 0.002$), which remained unchanged after adjusting for age, gender, LVEF, CAD, diabetes, hypertension, and valve surgery. The independent beneficial effects of beta-blockers were seen in patients with CAD and no CAD and in patients managed both surgically and conservatively.

The outcome of 1,344 patients operated on between 1980 and 1995 for severe MR was analyzed by age: < 65 years ($n = 556$), 65 to 74 years ($n = 504$), and ≥ 75 years ($n = 284$). At baseline, prevalence of severe symptoms, AF, CAD, "ischemic" MR, need for CABG, and creatinine level were

higher in older versus younger patients (all $p < 0.002$) (13). Over the time periods of 1980 to 1983, 1984 to 1987, 1988 to 1991, and 1992 to 1995, operative mortality fell progressively, in the two older subgroups from about 23% to 27% to about 5%, the incidence of low-output state decreased from about 38% to 50% to $\geq 10\%$ and the probability of mitral valve repair increased from approximately 30% to about 80% to 82% (all $p \leq 0.001$).

In patients with dilated cardiomyopathy, restrictive mitral annuloplasty (RMA) “effectively” restored MR without significant acute changes in baseline hemodynamics and left ventricular systolic function (14).

The EVEREST phase I clinical trial Evalve clip for percutaneous edge-to-edge mitral valve repair was performed in 20 patients, mean age 68 years, of which 75% were men (15). A partial clip detached in one patient, and surgery for MR was performed in two patients. Patients were discharged to home at an average time of 1.7 days. At 30 days’ follow-up, the results were “encouraging.” Posterior myocardial infarction (MI) was produced in six sheep and was compared to six control sheep. The percutaneous transvenous annuloplasty device was evaluated (16). The MR jet area, mitral valve tenting heights, and tenting area in all three commissures were significantly greater in chronic ischemic MR than in controls and were significantly reduced after coronary sinus-based annuloplasty.

TRICUSPID REGURGITATION (TR)

A total of 216 patients with functional TR were studied before and after tricuspid annuloplasty (17). Residual moderate TR was present in 15% and severe TR in 7.4% of patients. Multivariate analysis showed that age, tethering distance, and severity of preoperative TR were predictive of residual TR.

Severe symptomatic TR was associated with previous permanent pacemaker (PPM) or implantable cardioverter-defibrillator (ICD) (18). Forty-one patients with a mean age of 70 years (20 men; 21 women) had a normal tricuspid valve (TV) apparatus at surgery and 14 had dilated TV. The PPM lead TV perforation was found in 7, entanglement in TV leaflet or chordae in 4, impingement of the TV leaflets in 16 patients; adherence to the valve was seen in 14 patients. These findings were suspected in only 5 of 41 patients by preoperative transthoracic echocardiography and 4 of 13 patients in preoperative transesophageal echocardiography.

A total of 193 patients underwent MVR for rheumatic valve disease, of whom 57 also underwent tricuspid annuloplasty (TAP) (19). Of 21 patients (10.9%) who developed TR, mitral prosthetic valve function was normal. Severe TR developed in 11 of 35 (28.2%) patients who initially had severe TR. Despite successful TAP, the severity of TR at time of mitral valve surgery was the most important factor for prediction of severe TR after surgery.

Tricuspid valve replacement provided more complete

elimination of TR than did annuloplasty (20). Right ventricular function was reduced after TVR, whereas it was preserved after annuloplasty.

Fifty-two patients underwent TV repair for functional TR (grade ≥ 3) due to annular dilation with three-dimensional ring annuloplasty for TR (Edwards MC³ system) (21). At six-month follow-up, TR had a mean grade of 1.5 (range 0 to 2).

INFECTIVE ENDOCARDITIS (IE)

There was an unchanged pattern of IE. Over a period of 30 years (1970 to 2000), the population of Olmsted County, Minnesota, which consists largely of middle-class whites with an “extremely” low prevalence of intravenous (IV) drug abuse, was evaluated (22). The incidence of IE was relatively unchanged, and the leading causative organism was *Streptococcus viridans*.

In a multicenter, international study of prosthetic valve endocarditis (PVE), 355 patients were studied. Surgical treatment was utilized in 148 patients (41.6%), median time to surgery was 12 days, and 207 patients (59.4%) were treated medically (23). In-hospital mortality in the two groups was 24.3% and 22.7%, respectively ($p = 0.729$), and was “strongly related” to *S. aureus* infection and to the occurrence of cerebrovascular events with a trend toward improved survival with surgery. Surgery for complicated PVE resulted in a similar in-hospital survival rate as for uncomplicated PVE without surgery.

SURGERY

Postoperative AF occurred in 673 of 2,795 patients (24%) undergoing CABG (24). The incidence was higher in those with hypertension, chronic lung disease, cancer, and in subjects in New York Heart Association (NYHA) functional classes III/IV. Those who developed AF had longer stays in the intensive care unit and in the hospital, and also had a higher operative mortality (1.6% vs. 0.7%).

A prospective, randomized trial of four days of IV amiodarone was performed in 120 patients undergoing cardiac valve surgery (25). At four days, the incidence of AF was significantly lower in the amiodarone group (10% vs. 26%), but after drug discontinuation it was higher (37% vs. 19%).

A meta-analysis of 12 randomized trials of cardiothoracic surgery showed prophylactic amiodarone reduced the incidence of postoperative AF with an odds ratio (OR) of 0.47 (95% CI 0.39 to 0.57) but the differences of the effects on stroke and total hospital costs were not significantly different (26).

A prospective study of 648 patients undergoing cardiac surgery was performed at 14 VA Medical Centers (27). At baseline, 29.2% (189 of 648 patients) were depressed; they were younger; more were in NYHA functional classes III/IV, were on preoperative IV nitroglycerin or intra-aortic balloon pump, and had emergent surgery. Six-month mor-

tality was higher, 13.2% vs. 7.6%, in non-depressed patients ($p = 0.03$). On multivariable analysis, depressed patients had a higher mortality (OR 1.9, 95% CI 1.07 to 3.4; $p = 0.03$).

The 2001 Euro Heart Survey included 5,001 patients; 1,231 of these patients underwent valve surgery (28). The average age of the patients was 64 ± 13 years, with 43% in NYHA functional class I or II. The operative mortality for single-valve disease was 2.9%, for multiple-valve disease it was 6.5%, and for redo surgery it was 5.8%. The predictors of operative mortality were Euroscore ($p < 0.0001$), NYHA functional class ($p = 0.018$), and multiple versus single-valve surgery ($p = 0.021$).

A cohort of 240 consecutive asymptomatic or mildly symptomatic patients underwent aortic valve replacement (AVR) \pm CABG between 1992 and 2004 (29). The operative mortality for isolated AVR (AS, $n = 35$; aortic regurgitation [AR], $n = 82$) was 0%; for AVR + CABG for AS ($n = 9$) it was 0%; and AR ($n = 114$) it was 1.6%.

In a randomized trial of surgical correction of permanent AF, 51 patients were randomized either to no cryoablation (control), Cox-Maze III surgery, or isolation of pulmonary veins (IPV) (30). At follow-up of 28.4 ± 14 months, the incidence of AF in the control group was 56.3%, whereas incidence of sinus rhythm in the IPV group was 88.9% and in the Maze group it was 84.6%.

Patients who had percutaneous coronary intervention followed by AVR (hybrid procedures; $n = 20$) versus CABG + AVR (standard; $n = 114$) had a statistically significant higher percentage of women, those who had previous cardiothoracic surgery, pulmonary hypertension, and peripheral vascular disease (31). The 90-day mortality was similar (5% vs. 4%); 1-year mortality was also similar (6% vs. 9%) as was 1-year MACE incidence was also similar (11% vs. 9%).

PROSTHETIC HEART VALVES

In a prospective study of 89 consecutive patients, mean age 67 ± 13 years, who had prosthetic heart valves, 77.5% of 69 ischemic cerebral events were shown by cranial computed tomography to be due to bleeding. The international normalized ratio was >5.0 in seven patients and <1.8 in four patients (32).

Of 131 explants for structural valve deterioration (SVD) in bioprosthetic MVR, the only predictor for SVD was younger age ($p < 0.001$) (33). Total cholesterol, high-density lipoprotein and low-density lipoprotein cholesterol, CAD, and creatinine levels were not predictors.

Finally, 15 porcine valves of similar annulus size were studied in a pulse duplicator at various flow rates to determine effect of increasing flow on bioprostheses of different stiffness (34). As flow rates were increased from 2.0 to 5.75 l/min, peak pressure gradient increased from 16 to 73 mm Hg ($p < 0.003$), the aortic valve area (AVA) ratio increased from 1.16 to 2.10 ($p < 0.003$). However, AVA at

2.0 to 5.75 l/min was 0.94 versus 0.99 ($p = \text{NS}$) and AVA at different flow rates ranged from 0.85 to 0.99.

Reprint requests and correspondence: Dr. Shahbudin H. Rahimtoola, Distinguished Professor, University of Southern California, 2025 Zonal Avenue, Los Angeles, California 90033. E-mail: rahimtoo@usc.edu.

REFERENCES

- Novaro GM, Aviles RJ, Katz R, et al. Demographic characteristics, but not C-reactive protein, predict progression of calcific aortic valve disease (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:357A.
- Peters CM, Popovic Z, Novaro G, Stewart W. Are markers of atherosclerotic risk predictive of native aortic valve stenosis progression? A large retrospective study (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:357A.
- Skowasch D, Steinmetz M, Schrempf S, et al. Cellular characteristics of aortic valve degeneration. Endothelial progenitor-, dendritic-, and inflammatory-cell markers in degenerative aortic valve stenosis and explanted bioprostheses (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:358A.
- Lung B, Messika-Zeitoun D, Baron G, et al. What patient characteristics lead to contra-indication of surgery in elderly patients with severe symptomatic, aortic stenosis (abstr)? *J Am Coll Cardiol* 2005;45 Suppl A:362A.
- Eltchaninoff H, Bauer F, Tron C, Agatiello C, Nercolini D, Criber A. The I-REVIVE study: mid-term follow-up after percutaneous aortic valve implantation in non-surgical patients with calcific aortic stenosis (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:358A.
- Grube E, Laborde J-C, Seth A, Lal P, Bonan R. First human cases of percutaneous aortic valve replacement with the REVALVING system (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:358A.
- Robertson KA, Volmink JA, Mayosi BM. Evidence from a meta-analysis of randomized controlled trials shows that primary prevention of acute rheumatic fever with antibiotics is cost-effective in developing countries (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:355A.
- Ramondo A, Napodano M, Tarantini G, et al. Age-related outcome of percutaneous mitral balloon valvuloplasty (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:355A.
- Fawzy ME, Hassan W, Stefadouros M, Hegazy H, El Shaer F, Chaudhary M. Long-term results (up to 14 years) of mitral balloon valvotomy in a series of 468 patients and predictors of long-term outcome (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:356A.
- Pandya S, Subramaniam M, Stock S, et al. Degenerative versus rheumatic mitral valve disease: evaluation of cellular proliferation and osteoblast mediated extracellular matrix production (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:364A.
- Rosenhek R, Rader F, Gabriel H, et al. Surgery for severe but asymptomatic mitral regurgitation (abstr)? *J Am Coll Cardiol* 2005;45 Suppl A:363A.
- Varadarajan P, Appel D, Joshni N, Duvuri L, Pai RG. Protective effect of beta blocker therapy in chronic severe mitral regurgitation patients with normal ejection fraction (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:8A.
- Detaint D, Sundt T, Nkomo V, et al. Surgical correction of mitral regurgitation in the elderly: risks and recent improvements (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:14A.
- Tulner SA, Steendijk P, Klautz R, et al. Acute effects of restrictive mitral annuloplasty in patients with chronic heart failure (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:353A.
- Feldman T, Wasserman HS, Hermann HC, et al. Percutaneous edge-to-edge mitral valve repair using the Evalve clip: update of the EVEREST phase I clinical trial (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:359A.
- Daimon M, Shiota T, Liddicoat J, et al. Percutaneous mitral valve repair for chronic ischemic mitral regurgitation: real-time 3D echocardiographic study (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:1A.
- Moon JY, Shim CY, Ha J-W, et al. Tricuspid regurgitation late after mitral valve replacement for rheumatic mitral valve disease (abstr). *J Am Coll Cardiol* 2005;45 Suppl A:356A.

18. Lin G, Nishimura R, Connolly H, Dearani J, Sundt T, Hayes D. Severe symptomatic tricuspid valve regurgitation due to permanent pacemaker leads: fact or fiction (abstr)? J Am Coll Cardiol 2005;45 Suppl A:362A.
19. Fukuda S, Song-JM, Gillinov AM, et al. Tricuspid valve tethering predicts residual regurgitation after tricuspid annuloplasty (abstr). J Am Coll Cardiol 2005;45 Suppl A:362A.
20. Fukuda S, Gillinov AM, Song J-M, et al. The effect of tricuspid valve surgery on right ventricular function in patients with functional tricuspid regurgitation (abstr). J Am Coll Cardiol 2005;45 Suppl A:365A.
21. Filsoufi F, Salzberg SP, von Harbou K, Aklog L, Goldman ME, Adams DH. Three dimensional ring annuloplasty: a significant advance for the treatment of tricuspid regurgitation (abstr). J Am Coll Cardiol 2005;45 Suppl A:366A.
22. Tleyjeh IM, Murad HS, Ghomrawi HM, et al. Changing pattern of infective endocarditis: a population study of Olmsted County, Minnesota (abstr). J Am Coll Cardiol 2005;45 Suppl A:362A.
23. Wang A, Pappas P, Anstrom KJ, et al. The use and effect of surgical therapy for prosthetic valve infective endocarditis: a propensity analysis of a multicenter, international cohort (abstr). J Am Coll Cardiol 2005;45 Suppl A:363A.
24. Al-Khatib SM, Goyal AG, Hafley GE, et al. Short-term outcomes of atrial fibrillation complicating coronary artery bypass graft surgery: results from the PREVENT-IV trial (abstr). J Am Coll Cardiol 2005;45 Suppl A:365A.
25. Beaulieu Y, Denault AY, Couture P, et al. Randomized controlled trial on the use of intravenous amiodarone for prophylaxis of atrial fibrillation in patients undergoing cardiac valvular surgery (abstr). J Am Coll Cardiol 2005;45 Suppl A:352A.
26. Gillespie EL, Perkerson KA, White CM, Kluger J, Coleman CI. What is the impact of amiodarone on postoperative outcomes? A meta-analysis (abstr). J Am Coll Cardiol 2005;45 Suppl A:355A.
27. Ho PM, Masoudi FA, Spertus JA, et al. Depression predicts mortality following cardiac valve surgery (abstr). J Am Coll Cardiol 2005;45 Suppl A:355A.
28. Jung B, Baron G, Fondard O, et al. Is the Euroscore reliable in the prediction of operative mortality in valve surgery (abstr)? J Am Coll Cardiol 2005;45 Suppl A:357A.
29. Shekar PS, Aranki SF, Couper GS, Leacche M, Taft M, Cohn LH. What is the current operative risk of aortic valve surgery for asymptomatic aortic valve disease (abstr)? J Am Coll Cardiol 2005;45 Suppl A:365A.
30. Albrecht A, Kalil RA, de Lima GG, et al. Randomized study of surgical correction of permanent atrial fibrillation (abstr). J Am Coll Cardiol 2005;45 Suppl A:360A.
31. Meier MA, Deeb M, Chetcuti S, Moscucci M. Acute and long term outcomes of percutaneous coronary intervention prior to aortic root replacement (abstr). J Am Coll Cardiol 2005;45 Suppl A:351A.
32. Piper C, Hering D, Langer C, Horstkotte D. How to report morbid events in patients with prosthetic heart valves: a prospective study of patients presenting with cerebral events (abstr). J Am Coll Cardiol 2005;45 Suppl A:352A.
33. Gring C, Houghtaling P, Novaro G, et al. Pre-operative cholesterol levels do not predict structural valve degeneration in patients undergoing bioprosthetic mitral valve replacement (abstr). J Am Coll Cardiol 2005;45 Suppl A:360A.
34. Singh A, VanAuker MD, Rhodes K, Koob TJ, Strom JA. Effects of changes in cardiac output and leaflet properties on echocardiographic markers of aortic valve stiffness (abstr). J Am Coll Cardiol 2005;45 Suppl A:361A.